



СЕЧЕНОВСКИЙ
УНИВЕРСИТЕТ
НАУК О ЖИЗНИ



КФК



THIRD INTERNATIONAL CONFERENCE «INTEGRATION NETWORK OF THE PHARMACEUTICAL ECOLOGY - 2025»

EFFECT OF CYCLACHAENA XANTHIIFOLIA (NUTT.) FRESEN. EXTRACT ON THE GROWTH AND DEVELOPMENT OF THE ROOTS OF ALLIUM CEPA L.

Anastasiya Hryshan student, Lyubov Fedorova, senior lecturer

Pharmaceutical Natural Sciences of the A.P. Nelyubin Institute of Pharmacy, Sechenov University



Fig. 1 Cyclachena - object of study



Fig. 5 Results of the growth of the onion on a high concentration cyclachene with addition of potassium bichromate



Fig. 6 Results of the growth of onion on low concentration cyclachene with addition of potassium bichromate

Introduction

Cyclachena xanthiifolia (Nutt.) Fresen is an invasive annual plant with pronounced phytotoxicity affecting the ecosystem and human health. Evaluation of toxicity of plant extracts is convenient by the Allium-test method.

Results

- Cyclachene extracts caused mitotic delay and suppression of root formation: average root length 4-6 mm.
- The option «extract (more concentr.) + K₂Cr₂O₇» demonstrated partial removal of inhibition (7 mm) compared to the same extract without bichromate.
- Control (water): active root formation, 27 mm.
- Water + K₂Cr₂O₇ 0.4 mg/l: growth stimulation, 44 mm.
- The average inhibition rate of extracts relative to control is about 81%.



Fig. 7 Low concentration onion growth results



Fig. 8 High concentration onion growth results

Purpose

To evaluate the phytotoxicity of water extractions cyclenoses by the Allium-test method.

Materials and methods

Product: Allium cepa L., «Stütgarten rizes» class, 10-15 g.

Scheme of experience: in a container by 80 ml of solution; by 3 bulbs per option; triple repetition.

Medium: distilled water (control); water + K₂Cr₂O₇ 0.4 mg/l; extracts cyclene 1:20 and 1:10; their combinations with K₂Cr₂O₇.

Incubation: 7 days at 22-24 °C.

Estimate: length and condition of the roots.



Fig. 4 Results of growing onion on distilled water with bichromate




Fig. 3 Results of growing onion in distilled water



Fig. 2 Production of allium-test - growing onions in the studied environments

Conclusions

- The extracts of Cyclachena xanthiifolia in both concentrations have a pronounced phytotoxic effect on Allium cepa roots; the strength of the effect depends on the concentration.
- The addition of K₂Cr₂O₇ (0.4 mg/l) has a modifying effect: it antagonistically weakens the inhibition by the high concentration extract and itself causes growth stimulation in water.
- Allium-test is an indicator for rapid screening assessment of phytotoxicity of invasive species.




Fig. 9 Average length of the roots of the onion